



# Ecological Restoration Institute



Fact Sheet: Planting to Restore Ponderosa Pine Sites Burned by High-Severity Fire

September 2014

## Planting to Restore Ponderosa Pine Sites Burned by High-Severity Fire

By John Paul Roccaforte, ERI Research Specialist, Sr.

### INTRODUCTION

Increases in landscape-scale wildfires in frequent-fire forests over the last several decades have led to management concerns regarding long-term restoration of severely burned sites. In particular, interior areas of large, high-severity patches may lack conifer regeneration for decades (Savage and Mast 2005, Passovoy and Fulé 2006, Haire and McGarigal 2010, Roccaforte et al. 2012). A lack of tree regeneration may result in type conversion from sites historically dominated by coniferous forests to persistent non-forested areas (Figure 1) (Barton 2002, Strom and Fulé 2007). Based on these concerns, managers have at times aggressively pursued replanting of burned-over sites with conifers, with the result that many planted areas eventually become overly dense with trees to the point that eventually crown fires are bound to recur. To avoid this problem and to assure long-term restoration of forest structure including natural openings, consideration should be given to reestablishment of reference species composition, natural ranges and variability in tree densities, and spatial arrangement of trees in a functional ecosystem.

This fact sheet outlines a relatively new approach—one based on ecological restoration principles—for tree planting after high-severity wildfire in southwestern frequent-fire forests. Historically, land managers were directed to reestablish full site stocking with an emphasis on future timber production. Presently, concern for restoration of long-term resilience and ecological function directs managers to consider natural ranges of variability (reference conditions) in tree density and spatial arrangement to guide replanting strategies. We use the Forest Vegetation Simulator (FVS) to demonstrate planting densities that emulate pre-fire exclusion reference conditions at the mid-scale range of 10–1,000 acres (Reynolds et al. 2013). In the absence of subsequent management, higher initial planting densities will likely lead to poor forest health and increased crown fire hazard.



**Figure 1.** The Pot Fire (top) burned in 1996 and had no regeneration 11 years following the fire; the Aspen Fire (bottom) in southern Arizona burned in 2003 and has the potential of transitioning into an oak shrubfield.

The Ecological Restoration Institute is dedicated to the restoration of fire-adapted forests and woodlands. ERI provides services that support the social and economic vitality of communities that depend on forests and the natural resources and ecosystem services they provide. Our efforts focus on science-based research of ecological and socio-economic issues related to restoration as well as support for on-the-ground treatments, outreach and education.

Ecological Restoration Institute, P.O. Box 15017, Flagstaff, AZ 86011, 928.523.7182, FAX 928.523.0296, [www.nau.edu/eri](http://www.nau.edu/eri)

## **HISTORY OF PONDEROSA PINE PLANTING IN THE SOUTHWEST**

There is a rich history of research related to tree regeneration in southwestern forests dating back to the early 20th century. The Fort Valley Experiment Station was established in 1908 to study the apparent lack of sufficient natural regeneration in ponderosa pine forests (Fletcher 1979). Research from 1912 into the 1930s helped develop detailed guidelines for planting to achieve optimal seedling survival and stocking (Larson 1957, Hayes 1961; Heidmann 1963, see also Heidmann 2008). Schubert and his colleagues (1969) summarized these recommendations and produced a simple planting guide. Although guidelines concerning planting density are presently flexible and allow silviculturists to customize planting programs to meet local objectives (USFS Region 3 2002), managers historically aimed to reestablish full site stocking. For example, Schubert (1974) recommended planting 680 trees per acre at an 8-by-8-foot spacing based on the assumption that 50 percent of the seedlings would survive and attain a diameter of five inches. These 340 surviving, 5-inch trees would be needed for a growing stock level of 80 (Schubert 1974). Studies that discussed planting arrangement always recommended even spacing insofar as possible (e.g., Heidmann et al. 1977), although Schubert (1974) noted the importance of planting in the best spots, even at the expense of consistent spacing between trees.

Funding for reforestation programs has declined in recent decades (Burch 2005), although reforestation is still mandated under the Anderson-Mansfield Reforestation and Revegetation Joint Resolution Act of 1949 (at 16 U.S.C. 581j and 581 j(note)) and the National Forest Management Act of 1976.

## **USING REFERENCE CONDITIONS TO GUIDE PLANTING DECISIONS**

Extensive research over the last few decades has indicated that tree density in southwestern ponderosa pine forests historically ranged about 20 to 50 ponderosa pines per acre prior to the widespread interruption of the frequent surface fire regime (Stoddard 2011). From a timber production standpoint, sites were understocked (Woolsey 1911). In addition, before fire exclusion, trees often showed aggregated spatial patterns, with tree groups of varying sizes and single trees surrounded by a predominant matrix of grassy openings (Sánchez Meador et al. 2011, Larson and Churchill 2012). Tree groups ranged from less than one-tenth of an acre in size to nearly one acre, with two to 72 trees per group. Frequent surface fires and grass competition sufficiently maintained open conditions with regeneration rates as low as 1.5 trees per acre per decade (Mast et al. 1999, Bailey and Covington 2002). Open forests with scattered groups of trees provided a wide range of habitats for a diversity of animals and plant life, and were resilient to frequent occurrence of fire. Today, many of these forests are overly dense with trees due to decades of fire exclusion and are degraded in terms of forest health and resilience to wildfire (Covington and Moore 1994, Dahms and Geils 1997).

Replanting severely burned areas in a pattern that emulates natural patterns of healthy forests that historically occurred on the site is one way to promote resilient forests that function within their historical range of variability. For southwestern ponderosa pine forests, this translates to planting fewer trees than the traditional full stocking approach. On many sites, this also requires planting trees in various sized groups and leaving unplanted openings, which often occupied 60 to 80 percent of the area in presettlement stands (Covington and Sackett 1992, Covington and Moore 1994). In Table 1, Forest Vegetation Simulator (FVS) projections using varying planting densities and assumed survival rates suggest that multi-resource goals are likely to be met at intermediate planting levels. Table 1 shows that, depending on site index (the potential productive quality of an area where trees grow) and assuming 50 percent survival (Schubert 1974, Ouzts 2014), planting from 60–100 seedlings per acre will, after 80 years, result in tree densities similar to those found in historical tree density studies (Stoddard 2011) and recommended by Covington et al. (1997) for restoring stands to within natural ranges of structural variability. In contrast, FVS predictions indicate that planting 300 seedlings per acre will result in tree densities well above the historical range of variability reported by Stoddard (2011). Managers should plan post-fire reforestation efforts that promote resilient future forests while recognizing that stands planted with too many seedlings have the potential to become overly dense and susceptible to crown fire.

Although FVS does not account for variations in tree spatial arrangement, use of this tool can help land managers evaluate relative differences among planting scenarios. By studying these differences, practitioners can determine strategies for reaching forest health goals over time.

## FUTURE RESEARCH AND MONITORING

Monitoring and evaluation of successes and failures is needed for restoration of severe wildfire sites. Near-term monitoring should document the planting objectives and strategy, site selection criteria and site conditions, planting technique, density, and arrangement in addition to initial seedling survival and planting costs. Longer-term studies should continue to monitor tree survival but also assess responses of other important ecosystem components. An additional potential study could test drought-resistant genotypes in anticipation of climate change. Long-term monitoring will help managers strengthen adaptive management frameworks and ensure that sound, evidence-based decisions are made.

Seedling Survival = 50%													
Site Index = 70													
Year	Planting Density = 60 TPA				Planting Density = 100 TPA				Planting Density = 300 TPA				
	TPA	BA	SDI	Crown- ing	TPA	BA	SDI	Crown- ing	TPA	BA	SDI	Crown- ing	
1	30	0	0	0	50	0	0	0	150	0	0	0	
11	29	0	1	364	49	0	2	255	147	1	5	122	
21	29	2	6	169	48	4	10	119	144	11	31	57	
31	28	6	13	119	47	10	22	85	142	27	60	41	
41	28	11	22	102	46	18	35	73	139	43	88	37	
51	27	17	30	96	45	26	47	71	136	60	114	36	
61	27	22	38	94	45	34	58	69	134	75	137	36	
71	26	28	45	94	44	41	68	69	131	90	158	37	
81	26	33	51	95	43	48	77	69	129	104	177	38	
Site Index = 80													
Year	Planting Density = 60 TPA				Planting Density = 100 TPA				Planting Density = 300 TPA				
	TPA	BA	SDI	Crown- ing	TPA	BA	SDI	Crown- ing	TPA	BA	SDI	Crown- ing	
1	30	0	0	0	50	0	0	0	150	0	0	0	
11	29	1	2	314	49	1	3	222	147	3	10	105	
21	29	3	8	153	48	6	14	108	144	17	42	49	
31	28	8	17	115	47	14	28	81	142	35	75	39	
41	28	14	26	103	46	23	42	74	139	53	105	36	
51	27	20	35	99	45	32	55	71	136	72	132	36	
61	27	26	43	97	45	41	67	71	134	89	157	36	
71	26	32	50	98	44	49	78	72	131	106	179	37	
81	26	38	57	99	43	57	87	72	128	121	199	38	

**Table 1.** Forest Vegetation Simulator output for ponderosa pine planting scenarios at 50 percent seedling survival. Two site index values (70 and 80) and three planting densities (60, 100, and 300 TPA) were simulated. The output shows estimates of tree density (TPA), basal area (BA (ft<sup>2</sup> per acre)), Reineke's (1933) stand density index (SDI), and crowning index (Crowning (mph)) for years 1–81 from time of planting.

## LITERATURE CITED

- Barton, A.M. 2002 Intense wildfire in southeastern Arizona: a transformation of a Madrean oak-pine forest to oak woodland. *Forest Ecology and Management* 165:205-212.
- Bailey, J.D. and W.W. Covington. 2002. Evaluating ponderosa pine regeneration rates following ecological restoration treatments in northern Arizona, USA. *Forest Ecology and Management* 155:271-278.
- Burch, F. 2005. Twenty years of nursery history—a Forest Service perspective. In R.K. Dumroese, L.E. Riley, and T.D. Landis, tech. cords. National proceedings: Forest and Conservation Nursery Associations—2004; 2004 July 12-15; Charleston, NC; and 2004 July 26-29; Medford, OR. Proc. RMRS-P-35. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Covington, W.W., P.Z. Fulé, M.M. Moore, S.C. Hart, T.E. Kolb, J.N. Mast, S.S. Sackett, and M.R. Wagner. 1997. Restoring ecosystem health in ponderosa pine forests of the Southwest. *Journal of Forestry* 95:23-29.
- Covington, W.W. and M.M. Moore. 1994. Postsettlement changes in natural fire regimes and forest structure: ecological restoration of old-growth ponderosa pine forests. *Journal of Sustainable Forestry* 2:153-181.
- Covington, W.W. and S.S. Sackett. 1992. Soil mineral nitrogen changes following prescribed burning in ponderosa pine. *Forest Ecology and Management* 54:175-191.
- Dahms, C.W. and B.W. Geils., tech. eds. 1997. An assessment of forest ecosystem health in the Southwest. General Technical Report RM-GTR-295. Fort Collins, CO. U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 97 p.
- Fletcher, R. 1979. Regeneration in the southwest—it can work! *Forestry Research West*, November: 9-12.
- Haire, S.L. and K. McGarigal. 2010. Effects of landscape patterns of fire severity on regeneration ponderosa pine forests (*Pinus ponderosa*) in New Mexico and Arizona, USA. *Landscape Ecology* 25:1055-1069.
- Hayes, G.L. 1961. Seeding and planting research at the Rocky Mountain Forest and Range Experiment Station. Forest Management Research Work Conference. October 12-13, 1961. Minneapolis, MN.
- Heidmann, L.J. 1963. Planting ponderosa pine successfully in the Southwest. Society of American Foresters Chapter Meeting.
- \_\_\_\_\_. 2008. Forest regeneration research at Fort Valley. Pp. 25-37 in S.D. Olberding and M.M. Moore, tech coords., Fort Valley Experimental Forest—A century of research 1908-2008. Proceedings RMRS-P-55. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Heidmann, L.J., F.R. Larson, and W.J. Rietveld. 1977. Evaluating ponderosa pine reforestation techniques in Central Arizona. USDA Forest Service Research Paper RM-190. Rocky Mountain Forest and Range Experiment Station. Ft. Collins, CO.
- Larson, M.M. 1957. Observations of 1-year-old ponderosa pine seedlings during a drought year in Arizona. Final Report.
- Larson, A.J. and D. Churchill. 2012. Tree spatial patterns in fire-frequent forests of western North America, including mechanisms of pattern formation and implications for designing fuel reduction and restoration treatments. *Forest Ecology and Management* 267:74-92.
- Mast, J.N., P.Z. Fulé, M.M. Moore, W.W. Covington, and A.E.M. Waltz. 1999. Restoration of presettlement age structure of an Arizona ponderosa pine forest. *Ecological Applications* 9:228-229.
- Ouzts, J.R. 2014. Post-fire ponderosa pine plantings in the southwest U.S.A. M.S. Thesis, Northern Arizona University, Flagstaff, AZ.
- Passovoy, M.D. and P.Z. Fulé. 2006. Snag and woody debris dynamics following severe wildfires in northern Arizona ponderosa pine forests. *Forest Ecology and Management* 223:237-246.
- Reineke, L.H. 1933. Perfecting a stand-density index for even-aged forests. *Journal of Agricultural Research* 46:627-638.
- Reynolds, R.T., A.J. Sánchez Meador, J.A. Youtz, T. Nicolet, M. Matonis, P.L. Jackson, D.G. DeLorenzo, and A.D. Graves. 2013. Restoring composition and structure in Southwestern frequent-fire forests: A science-based framework for improving ecosystem resiliency. Gen. Tech. Rep. RMRS-GTR-310. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 76 p.
- Roccaforte, J.P., P.Z. Fulé, W.W. Chancellor, and D.C. Laughlin. 2012. Woody debris and tree regeneration dynamics following severe wildfires in Arizona ponderosa pine forests. *Canadian Journal of Forest Research*. 42: 593-604.
- Sánchez Meador, A.J., P.F. Parysow, and M.M. Moore. 2011. A new method for delineating tree patches and assessing spatial reference conditions of ponderosa pine forests in northern Arizona. *Restoration Ecology* 19:490-499.
- Savage, M. and J.N. Mast. 2005. How resilient are southwestern ponderosa pine forests after crown fires? *Canadian Journal of Forest Research* 35:967-977.
- Schubert, G.H. 1974. Silviculture of southwestern ponderosa pine: The status of our knowledge. USDA Forest Service Research Paper RM-123. Rocky Mountain Forest and Range Experiment Station. Ft. Collins, CO.
- Schubert, G.H., R.W. Pearl, and L.J. Heidmann. 1969. Here's how—a guide to planting in the Southwest. USDA Forest Service Research Paper RM-49. Rocky Mountain Forest and Range Experiment Station. Ft. Collins, CO.
- Stoddard, M.T. 2011. *Compilation of historical forest structure characteristics across the Southern Colorado Plateau*. Ecological Restoration Institute Fact Sheet. Northern Arizona University, Flagstaff, AZ.
- Strom, B.A. and P.Z. Fulé. 2007. Pre-wildfire fuel treatments affect long-term ponderosa pine forest dynamics. *International Journal of Wildland Fire* 16: 128-138.
- U.S. Forest Service Region 3. 2002. Forest Service Handbook 2409.17. Silvicultural practices handbook, Chapter 2 Reforestation. Albuquerque: U.S. Forest Service Region 3 Office.
- Woolsey, T.S., Jr. 1911. Western yellow pine in Arizona and New Mexico. U.S. Department of Agriculture, Forest Service Bulletin 101. Government Printing Office, Washington DC. 64 p.

### Contact

John Paul Roccaforte, [John.Roccaforte@nau.edu](mailto:John.Roccaforte@nau.edu)

---

NAU is an equal opportunity provider.

This research was funded by a grant from the USDA Forest Service.